Journal of Surgical Oncology

Volume 76

January, February, March, April 2001

Editor-in-Chief

Walley J. Temple, MD, FRSCS, FACS

University of Calgary
Tom Baker Cancer Centre
Calgary, Alberta, Canada

Wiley-Liss, Inc., New York

Journal of Surgical Oncology

Table of Contents

Volume 76, Number 1, January 2001

Guest Editorial: Multivisceral Resections for Colorectal Cancer
Marvin J. Lopez 1

Factors Influencing Choice Between Mastectomy and Lumpectomy for Women in the Carolinas
Susan Benedict, David J. Cole, Lisa Baron, and Paul Baron 6
Characteristics of women choosing between mastectomy and lumpectomy were determined. Sources of treatment-related information for the women included popular media, friends, relatives, cancer organizations, and the Internet.

Highly Sensitive Immunohistological Study for Detection of Estrogen Receptor in Human Breast Cancer
Yoshiaki Sumiyoshi, Takayuki Shirakusa, Yuichi Yamashita, Takafumi Maekawa, Toshimi Sakai, and Masahiro Kikuchi 13
This new, highly sensitive immunohistological method is useful in formalin–fixed, paraffin-embedded tissue, and in small tumor specimen.

Rat Glioma Cell Death Induced by Cationic Liposome-Mediated Transfer of the Herpes Simplex Virus Thymidine Kinase Gene Followed by Ganciclovir Treatment
Tazuka Yoshida, Masaaki Mizuno, Katsumi Taniguchi, Norihisa Nakayashiki, Toshihiko Wakabayashi, and Jun Yoshida 19
Introduction into rat glioma cell cultures of unilamellar carionic liposomes containing the herpes simplex virus thymidine kinase gene, followed by exposure to ganciclovir, resulted in tumor cell apoptosis and necrosis. Injection of the DNA-liposome complexes into experimental rat gliomas, again followed by ganciclovir administration, minimized tumor growth and increased host survival.

Expression of Cyclooxygenase-2 in Human Gastric Adenomas and Adenocarcinomas
Kazuhiko Uefuji, Takashi Ichikura, and Hidetaka Mochizuki 26
Cyclooxygenase (COX)-2 mRNA was detected in 19 of 37 (51%) human gastric cancer specimens by RT-PCR technique. COX-2 mRNA expression was associated with the size of the tumor, the depth of invasion, and lymph node involvement in gastric cancer, thus suggesting that COX-2 may play an important role in tumor progression of human gastric cancer.

Prognostic Significance of Expression of Thymidine Phosphorylase and Vascular Endothelial Growth Factor in Human Gastric Carcinoma
Hironobu Kimura, Kohji Konishi, Takefumi Nukui, Masahide Kaji, Kiichi Maeda, Kazuhisa Yabushita, Masahiko Tsuji, and Atsuo Miwa 31
Combination analysis of dThdPase and vascular endothelial growth factor expression in gastric carcinoma showed them to be well characterized inductors of prognosis and metastasis.

Clinicopathologic Study of Mixed Hepatocellular and Cholangiocellular Carcinoma: Modes of Spreading and Choice of Surgical Treatment by Reference to Macroscopic Type
Atsushi Sasaki, Katsunori Kawano, Masanori Aramaki, Tsuyoshi Ohno, Koichiro Tahara, Yu Takeuchi, Takanori Yoshida, and Seigo Kitano 37
Clinicopathologic features and modes of spreading of mixed hepatocellular and cholangiocellular carcinoma (MHC) differ according to macroscopic appearance. Lymphadenectomy might be appropriate in selected cases of MHCs.

Pulmonary Metastasectomy: Might the Type of Resection Affect Survival?
Tommaso Claudio Mineo, Vincenzo Ambrogi, Giuseppe Tonini, and Italo Nofroni 47
The impact on survival for different types of pulmonary metastasectomy was investigated in 85 patients. Conventional laser resections and lobectomy did not disclose statistically significant differences on survival, but the laser provided less morbidity and hospitalization.

Clinicopathological Characteristics of Surgically Resected Lung Cancer Associated With Idiopathic Pulmonary Fibrosis
Hidenori Kawasaki, Kanji Nagai, Tomoyuki Yokose, Junji Yoshida, Mitsuyo Nishimura, Kenro Takahashi, Kenji Suzuki, Ryutarou Kakinuma, and Yutaka Nishiwaki 53
Idiopathic pulmonary fibrosis (IPF) is well-known to be associated with lung cancer. To clarify clinicopathological features of IPF-related lung cancer, we compared lung cancer patients with and without IPF.

Reappraisal of the Role of NM23-H1 in Colorectal Cancers
Jenq-Chang Lee, Yih-Jyn Lin, Nan-Haw Chow, and Shan-Tair Wang 58
There was no apparent correlation between NM23-H1 expression and clinicopathological indicators, including liver metastasis. Determination of NM23-H1 expression didn't provide independent prognostic information compared with conventional pathological staging.

Expression of Fas Ligand is an Early Event in Colorectal Carcinogenesis
Masaaki Shimoyama, Tatsuo Kanda, Lili Liu, Yu Koyama, Takeyasu Suda, Yasuo Sakai, and Katsuyoshi Hatakeyama 63
Via immunohistochemical analysis of surgically resected colorectal specimens, the authors demonstrate that Fas ligand (FasL) was expressed in 81% of adenocarcinomas, 64% of adenomas with high-grade atypia, and 36% of adenomas with low-grade atypia. These results suggest that FasL expression may be a pivotal event in colorectal carcinogenesis.

Commentary
Edward R. Sauter 69

How I Do It: Fertility Preserving Testicular Transposition in Patients Undergoing Inguino Pelvic Irradiation
S.V.S. Deo, Sonal Asthana, Nootan K. Shukla, and Brigu K. Mohanti 70

Review Article: Interstitial Laser Thermotherapy in the Treatment of Colorectal Liver Metastases
Vijayaragavan Muralidharan and Christopher Christophi 73

Volume 76, Number 2, February 2001

Correlation of MTS1/p16 and nm23 mRNA Expression With Survival in Patients With Peripheral Synovial Sarcoma
Rastko Golouh, Giorgio Stanta, Matej Bračko, and Serena Bonin 83
In a series of 31 patients with synovial sarcoma who were followed for a median of 83 months, the presence of extensive tumor necrosis and high levels of nm23 mRNA expression were the only independent factors predicting survival.

Splenectomy in Proximal Gastric Cancer: Frequency of Lymph Node Metastasis to the Splenic Hilus
Stefan P. Mönig, Peter H. Collet, Stephan E. Baldus, Katrin Schmackpfeffer, Wolfgang Schröder, Jürgen Thiele, Hans P. Dienes, and Arnulf H. Hölscher 89
Lymph node metastasis to the splenic hilus was found in advanced proximal gastric cancer (UICC IIIb/IV) located at the greater curvature and in Borrmann type III/IV cancer with advanced lymph node metastasis.

Commentary
Walter Lawrence, Jr. 93

Combination Technique Is Superior to Dye Alone in Identification of the Sentinel Node in Breast Cancer Patients
Kazuyoshi Motomura, Hideo Inaji, Yoshifumi Komoike, Yoshihisa Hasegawa, Tsutomu Kasugai, Shinzaburo Noguchi, and Hiroki Koyama 95
The dye and radioisotope techniques are complementary. The combination of dye and radioisotope improve the detection rate of sentinel nodes and the diagnostic accuracy of axillary lymph node status over dye alone in patients with breast cancer.

Statistical Analysis of Diagnostic Failure of Fine Needle Aspiration Cytology (FNAC) in Breast Cancer
Kanji Tanaka, Tetsuji Shoji, Yuji Tominaga, Hiroyuki Nakagawa, Daigo Yamamoto, Hiroshi Kawanishi, Noriko Sakaida, Akiharu Okamura, and Koshiro Hioki 100
This study is aimed to clarify the interrelationship between the false-negative result of fine needle aspiration cytology on breast cancer and clinicopathological factors of patients with the use of uni- and multivariate analysis.

Pharmacokinetics of Intraperitoneal Oxaliplatin: Experimental Studies
Sophie R. Pestieau, Jim F. Belliveau, Hank Griffin, O. Anthony Stuart, and Paul H. Sugarbaker 106
Oxaliplatin was shown to be a pharmacokinetically advantageous drug for intraperitoneal administration; the addition of hyperthermia caused a trend toward increased oxaliplatin tissue concentration. Due to the likelihood of loco-regional recurrence after resection of a colon cancer, heated intraoperative intraperitoneal oxaliplatin may be suggested to prevent these recurrences.

Expression of CD44 Isoforms in Squamous Cell Carcinoma of the Border of the Tongue: A Correlation With Histological Grade, Pattern of Stromal Invasion, and Cell Differentiation
Isabel Fonseca, Teresa Pereira, Jorge Rosa-Santos, and Jorge Soares 115
Immunoexpression of CD44 variant isoforms was studied in a series of 56 cases of squamous cell carcinomas of the tongue and the results correlated with their histological grade, cell differentiation, and pattern of stromal invasion. Down-regulation of CD44v was found to relate to tumor grade, pattern of stromal invasion, and pN status.

Infrequent Microsatellite Instability in Biliary Tract Cancer
Takayuki Suto, Wataru Habano, Tamotsu Sugai, Noriyuki Uesugi, Senji Kanno, Kazuyoshi Saito, and Shin-Ichi Nakamura 121
We investigated microsatellite instability (MSI) and allelic loss at the hMLH1 and hMSH2 gene loci in 38 cases of extrahepatic bile duct (EHBD) cancer and 16 cases of ampullary cancer using polymerase chain reaction and an automated fluorescent DNA sequencer. Our results suggest that (a) genetic defects in the DNA mismatch repair system and MSI do not play an important role in the majority of biliary tract cancers and (b) MSI is not an independent predictor of prognosis in EHBD cancer.

Repeat Hepatectomy for Colorectal Liver Metastases: A Worthwhile Operation?
Andrea Muratore, Roberto Polastri, Hedayat Bouzari, Vincenzo Vergara, Alessandro Ferrero, and Lorenzo Capussotti 127
Repeat hepatic resection for colorectal metastases can provide long-term survival. A resectable synchronous extrahepatic disease is not a contraindication to surgery.

Prognostic Value of the Colposcopic Tumor Size in Stage IB Squamous Cervical Cancer
Kenji Yanoh, Nobuhiro Takeshima, Hidetaka Nishida, Yasuo Hirai, Nagayasu Toyoda, and Katsuhiko Hasumi 133
The colposcopic tumor size in stage IB squamous cervical cancer is an independent prognostic factor that can be used as an indicator for treatment options.

How I do It: Safe Pancreaticojejunostomy After Whipple Procedure: Modified Technique
Juan Celis, Francisco Berrospi, Eloy Ruiz, and Eduardo Payet 138

How I Do It: Simple Technique of Early Identification of the Thoracodorsal Nerve During Axillary Dissection
Vijay P. Khatri, Thelma Hurd, and Stephen B. Edge 141

Commentary
Constantine Karakousis 143

Review Article: Sentinel Lymph Node Biopsy as an Alternative to Routine Axillary Lymph Node Dissection in Breast Cancer Patients
Masakuni Noguchi 144

Volume 76, Number 3, March 2001

Fibrous Dysplasia of Bone: Management and Outcome of 20 Cases
Lucien C.M. Keijser, Tony G. Van Tienen, H.W. Bart Schreuder, Josef A.M. Lemmens, Maciej Pruszczynski, and Renè P.H. Veth 157
In a series of 20 patients with fibrous dysplasia of bone, the management and functional outcome of 32 fibrous lesions is reported. The functional outcome was uniformly good for patients with monostotic or polyostotic disease but unsatisfactory for patients with McCune-Albright syndrome.

Commentary
Timothy A. Damron 167

Activation of Matrix Metalloproteinase-2 is Correlated With Invasiveness in Thymic Epithelial Tumors
Kazuya Kondo, Hidetaka Kinoshita, Hisashi Ishikura, Takanori Miyoshi, Toshiyuki Hirose, Yasumichi Matsumori, and Yasumasa Monden 169
We examined the gelatinolytic activity of matrix metalloproteinase (MMP)-9, and the activity of active and inactive forms of MMP-2 in five thymi, five noninvasive thymomas, eight invasive thymomas, and five thymic carcinomas by quantitative gelatinolytic zymography. The gelatinolytic activity of active MMP-2 significantly correlated with the invasiveness in thymic epithelial tumors.

Clinical Significance of E-Cadherin Expression in Thyroid Neoplasms
Akihiro Naito, Hirotaka Iwase, Tatsuya Kuzushima, Takaaki Nakamura, and Shunzo Kobayashi 176
E-cadherin expression was measured in normal thyroid tissue and in thyroid adenomas and carcinomas. A loss of E-cadherin expression may be involved in regional lymph node metastasis and in malignant potential of thyroid neoplasms.

Instability of Chromosome 8 as an Indicator of Aggressive Tumor Phenotype in Pancreatic Cancer
Norihiro Sato, Kazuhiro Mizumoto, Naoki Maehara, Masaki Shono, Eishi Nagai, and Masao Tanaka 181
Chromosome instability is a common feature of pancreatic carcinoma. We show here that numerical variations of chromosome 8 correlate with aggressive tumor phenotype of pancreatic cancer cells.

Expression of Bcl-2 and p53 Protein in Resectable Invasive Ductal Carcinoma of the Pancreas: Effects on Clinical Outcome and Efficacy of Adjuvant Chemotherapy
Yoshinori Nio, Ming Dong, Chikage Iguchi, Kunihiro Yamasawa, Tomoko Toga, Masayuki Itakura, and Katsuhiro Tamura 188
In resectable invasive ductal carcinoma of the pancreas, bcl-2 expression is a beneficial prognostic factor and p53 expression may be a beneficial predictor for the effects of adjuvant chemotherapy.

Case Report: Primary Kaposi Sarcoma of the Bowel in a HIV-Negative Patient
Luigi Insabato, Dolores Di Vizio, Luigi Maria Terracciano, Guido Pettinato, Stefania Staibano, Delfina Bifano, and Luciano D'Agostino 197

Review Article: Statistical Methods in Cancer Research
Pippa M. Simpson, John A. Spratt, and John S. Spratt 201

Review Article: Psychosocial Issues: Pelvic Exenterative Surgery
Danielle Turns 224

Letter to the Editor: Continent Gastrostomy
John A. Clark and Robyn A. Pugash 237

Letter to the Editor: Reply
Pericles P. Vassilopoulos and Nickos Kelessis 238

Letter to the Editor: Is Postirradiation Angiosarcoma of the Breast so Rare and Does Breast Lymphedema Contribute to Its Development?
Csaba Polgár, Zsolt Orosz, and János Fodor 239

Letter to the Editor: Reply Drs. Polgar, Orosz, and Fodor
James Majeski 241

Volume 76, Number 4, April 2001

Guest Editorial: Hepatic Colorectal Metastases and Extrahepatic Disease
Jean-Nicolas Vauthey 243

Relationship Between Excision Volume, Margin Status, and Tumor Size With the Development of Local Recurrence in Patients With Ductal Carcinoma-In-Situ Treated With Breast-Conserving Therapy
Frank A. Vicini, Larry L. Kestin, Neal S. Goldstein, Kathy L. Baglan, Jane E. Pettinga, and Alvaro A. Martinez 245
The interrelationship between excision volume, margin status, and tumor size with local recurrence was examined in 146 patients with DCIS treated with breast-conserving therapy. Multivariate analysis revealed that patient age, margin status, the number of slides containing DCIS, and the number of DCIS/cancerization of labules foci within 5 mm of the margin, and an excision volume < 60 cm^3 were all independently associated with local recurrence risk.

Is the UICC Pathological Node Status System Useful? Comparison With the Japanese Breast Cancer Society Pathological Node Status System
Mitsuhiro Hayashi, Goi Sakamoto, Futoshi Akiyama, Fujio Kasumi, and Katsuyoshi Hatakeyama 255
We compared pathological node status of the UICC classification (UICC-NS) with that of the Japanese Breast Cancer Society (JBCS-NS).

Paget Disease of the Breast—an Easily Overlooked Disease?
Shyr-Ming Sheen-Chen, Han-Shiang Chen, Wei-Jen Chen, Hock-Liew Eng, Chih-Wei Sheen, and Fong-Fu Chou 261
Paget disease of the breast without a palpable breast mass carries a more favorable prognosis. Patients with any nipple complaints deserve a detailed evaluation even in the absence of a palpable breast mass in order not to overlook a favorable disease.

The Prognostic Significance of Preoperative Serum CA 19-9 in Patients With Resectable Gastric Carcinoma: Comparison With CEA
Nüvit Duraker and Ali Naci Çelik 266
In resectable gastric carcinoma, preoperative serum CA 19-9 and carcinoembryonic antigen levels may indicate stage of the disease, but neither has an independent prognostic value.

Pathology and Prognosis of Mucinous Gastric Carcinoma
Kazuhiro Yasuda, Yosuke Adachi, Norio Shiraishi, Kimio Yamaguchi, Akio Shiromizu, and Seigo Kitano 272
Multivariate analysis showed that mucinous histologic type was not a significant prognostic factor in patients with gastric carcinoma, and survival rate after gastrectomy for mucinous gastric carcinoma was not different from that for nonmucimous gastric carcinoma, when compared in the same category of tumor size, depth of wall invasion, lymph node metastasis, and stage.

Prognosis of T4 Gastric Carcinoma Patients: An Appraisal of Aggressive Surgical Treatment
Dipok Kumar Dhar, Hirofumi Kubota, Mitsuo Tachibana, Shoichi Kinugasa, Reiko Masunaga, Muneaki Shibakita, Hitoshi Kohno, and Naofumi Nagasue 278
Results of this retrospective review indicate that patients with gastric cancer with invasion into the surrounding organs might benefit from an aggressive resection of the tumor and invaded organs.

Epidemiologic Differences Between Women With Colorectal Cancer and Women With Ovarian Cancer
Gamal H. Eltabbakh, Nachimuthu Natarajan, M. Steven Piver, and Curtis J. Mettlin 283
The epidemiologic features of women with colorectal cancer are different from those with ovarian cancer. The difference between both groups might indicate difference in the environmental or genetic etiology of both cancers.

Solid-Pseudopapillary Tumor of the Pancreas (Frantz tumor) in Children: Report of Four Cases and Review of the Literature
Winfried Rebhandl, Franz Xaver Felberbauer, Stefan Puig, Kurosh Paya, Silvia Hochschorner, Murat Barlan, and Ernst Horcher 289
An important differential diagnosis of pancreatic tumors especially in young females. Four new cases are described and clinical work-up, surgical treatment, and histological features are discussed.

How I Do It: Intraductal Cooling of the Main Bile Ducts During Intraoperative Radiofrequency Ablation
Elias Dominique, Azzedine El Otmany, Alain Goharin, David Attalah, and Thierry De Baere 297

Treatment of Neoplastic Diseases of the Sacrum
George H. Raque, Jr., Todd W. Vitaz, and Christopher B. Shields 301
This article reviews the anatomy, pathology, and clinical presentation for various types of sacral lesions. In addition, it also reviews the various surgical procedures and potential complications associated with these techniques.

Letter to the Editor: Mucin Production as a Prognostic Factor in Resected Pancreatic Carcinoma
Pierre-Edouard Queneau and Massimo Usel 308

Author Index to Volume 76 309

Subject Index to Volume 76 311